

Computer-assisted proof of a periodic solution in a non-linear feedback DDE

Mikołaj Zalewski

Institute of Mathematics, Jagiellonian University

Reymonta 5, 30-059 Kraków, Poland

E-mail: Mikolaj.Zalewski@im.uj.edu.pl

Abstract

In this paper we rigorously prove the existence of a non-trivial periodic orbit for the non-linear DDE: $x'(t) = K \sin(x(t-1))$ for $K = 1.6$. We show that the equations on the Fourier coefficients have a solution by computing the local Brouwer degree. This degree can be computed by using a homotopy whose validity can be checked by checking a finite number of inequalities. Checking these inequalities is done by a computer program.

1 Introduction

We will show that the equation:

$$x'(t) = -K \sin(x(t-1)) \quad (1)$$

has a periodic solution for $K = 1.6$. By a solution we mean a $\mathcal{C}^1$ function that satisfies the equation.

Numerical simulations suggest that for $\frac{\pi}{2} < K < 5.1$ there is an attracting periodic orbit oscillating around 0. An informal argument for the existence of such an orbit also comes from the analysis of eigenvalues of the linearization [Wischert et al 94]. We will prove rigorously that for $K = 1.6$ this orbit exists.

In the remainder of the introduction we briefly outline our method. We will use the Fourier coefficients of the periodic orbit. First we rescale the time to obtain a 2π period. We make a substitution: $\tilde{x}(t) = x\left(\frac{t}{\tau}\right)$ where τ is a parameter. If τ is equal to $\frac{2\pi}{T}$ (where T is the period of the solution in the original equation) we will have a 2π -periodic solution of the new equation:

$$\begin{aligned}\tilde{x}'(t) &= \frac{1}{\tau} x' \left(\frac{t}{\tau} \right) = -\frac{K}{\tau} \sin \left(x \left(\frac{t}{\tau} - 1 \right) \right) = \\ &= -\frac{K}{\tau} \sin \left(x \left(\frac{t - \tau}{\tau} \right) \right) = -\frac{K}{\tau} \sin(\tilde{x}(t - \tau))\end{aligned}$$

By renaming $\tilde{x}$ to x we obtain:

$$x'(t) = -\frac{K}{\tau} \sin(x(t - \tau)) \quad (2)$$

We will prove for this equation that there exist a τ from a small interval such there exist a 2π -periodic orbit in a small neighborhood of a specified function. Note that we don't obtain the exact value of the period (even if numerical solution suggests it is 4, i.e. $\tau = \frac{\pi}{2}$) but treat τ as a variable.

To prove the existence of the orbit we will use the method of self-consistent bounds that was introduced in the context of Kuramoto-Shivashinsky PDEs in [Zgliczyński 01], [Zgliczyński 04]. When applied to the boundary value problem for ODEs or DDEs this method is similar to the Cesari method introduced in [Cesari 64] but doesn't require one of the conditions – see Section 2.4 in [Zgliczyński 01] for a comparison. Below we briefly describe the method.

We will derive from (2) the equation on Fourier coefficients. Let's first write (2) with a Taylor expansion instead of sinus:

$$x'(t) = -\frac{K}{\tau} \sum_{n=0}^{\infty} \frac{(-1)^k}{(2k+1)!} [x(t - \tau)]^{2k+1}$$

The equations on the Fourier coefficients will look similar:

$$\forall n \in \mathbb{Z} : inc_n = -\frac{K}{\tau} \sum_{k=0}^{\infty} \frac{(-1)^k}{(2k+1)!} (c^{*(2k+1)})_n e^{-in\tau}$$

where $(c^{*(2k+1)})_n$ means convoluting the sequence c with itself $2k+1$ times and then taking the n -th coefficient. The operation of convolution and this notation is introduced in details in section 2.

We will show that such a function has a non-trivial zero:

$$F(\tau, c) = \left\{ in\tau e^{in\tau} c_n + K \sum_{k=0}^{\infty} \frac{(-1)^k}{(2k+1)!} (c^{*(2k+1)})_n \right\}_{n=-\infty}^{\infty}$$

Where the domain is:

$$X_\beta := \left\{ x : \mathbb{Z} \rightarrow \mathbb{C} \mid \forall n \in \mathbb{Z} : |x_n| \leq \frac{\beta}{(|n| + 1)^2} \text{ and } x_n = \overline{x_{-n}} \right\}$$

for some $\beta > 0$. On X_β we consider the product topology (which is equivalent to component-wise convergence). It turns out that on such a domain the sums in the definition of F are convergent, F is continuous and each sequence $c \in X_\beta$ corresponds to a real-valued continuous function. Moreover if $F(c) = 0$ then the function is $\mathcal{C}^1$.

We will search for a zero in the neighborhood of an approximated solution obtained from numerical simulations. We will denote by $\hat{c}$ this approximation. For $n \leq n \leq 5$ it is equal to:

n	$\hat{c}_n$
0	0
1	$-0.1521000000 - 0.1163508047i$
2	0
3	$0.0001123121 - 0.0002746107i$
4	0
5	$-0.0000008173 - 0.0000001014i$

For $n > 5$ the $\hat{c}_n$ is zero, for $n < 0$ we have $\hat{c}_n = \overline{\hat{c}_{-n}}$. The τ in the approximation is: $\hat{\tau} = 1.570796$.

For each $l \in \mathbb{N}$ let us define the Galerkin projection P_l and immersion Q_l :

$$P_l : \mathbb{C}^{\mathbb{Z}} \ni c \rightarrow (c_0, \dots, c_l) \in \mathbb{R} \times \mathbb{C}^l$$

$$Q_l : \mathbb{R} \times \mathbb{C}^l \ni (c_0, \dots, c_l) \rightarrow (\dots, 0, 0, \overline{c_{-l}}, \dots, c_0, \dots, c_l, 0, 0, \dots) \in \mathbb{C}^{\mathbb{Z}}$$

Let's note that we need only the non-negative terms in the finite space as the negative terms can be obtained by conjugation. Also as $c_0 = \overline{c_{-0}}$ so we have $c_0 \in \mathbb{R}$. From the compactness of X_β it will be easy to show that:

Lemma. 15 *Let $\beta > 0$, $l_0 > 0$, $\underline{\tau}, \overline{\tau} \in \mathbb{R}$ be fixed. If for each $l > l_0$ there is a $c^l \in X_\beta$ and $\tau_l \in [\underline{\tau}, \overline{\tau}]$ such that $P_l F(\tau_l, c^l) = 0$ then there exist a $(c^0, \tau_0) \in X_\beta \times [\underline{\tau}, \overline{\tau}]$ such that $F(\tau_0, c^0) = 0$.*

Moreover if all the c^l are in a closed set D then $c^0 \in D$.

Thus it's enough that there is a zero for every Galerkin projection of F :

$$\tilde{F}_l : \mathbb{R} \times (\mathbb{R} \times \mathbb{C}^l) \ni (\tau, c) \rightarrow P_l F(\tau, Q_l(c)) \in \mathbb{R} \times \mathbb{C}^l$$

This will allow us to use a finite-dimensional topological method to show a zero of F . Let's note that the real dimension of the domain is $2l + 2$

while of the image is $2l + 1$ so if there is a zero then one can expect a 1-dimensional manifold of zeros. This manifold can be easily identified – if $x(\cdot)$ is a solution then $x(\cdot + \phi)$ is also. This means that if $\tilde{F}_l(\tau, c_0, c_1, \dots, c_l) = 0$ then $\tilde{F}_l(\tau, c_0, e^{i\phi}c_1, \dots, e^{il\phi}c_l) = 0$ what can be easily checked. To use the topological method we want to have the zero isolated so we will limit the domain – we assume that $c_1 \in \hat{c}_1 + \mathbb{R}$. Of course finding a zero in a limited domain implies a zero of the full system. By F_l we will denote the $\tilde{F}_l$ with the smaller domain.

The before-mentioned method to prove the existence of a zero of F_l is the local Brouwer degree (introduced e.g. in [Smoller 83]). Let's denote by $\deg(F_l, U, x)$ the degree of x on U . It is known that if the degree is non-zero then $\exists y \in U : f(y) = x$. We will use a neighborhood of $\hat{c}$ as U and $x = 0$.

To compute the degree we will use the homotopy invariance of the local Brouwer degree. As our homotopy H_l we will use a linear deformation of F_l into a function G_l that contains the most important terms of F_l (that is not strictly a linearization of F_l but it is close to):

$$H_l(h, \tau, c) = hF_l(\tau, c) + (1 - h)G_l(\tau, c)$$

The degree of G_l will be easy to show that it is non-zero. What we will need to show is that $0 \notin H_l([0; 1]; \partial U)$. To show that it's enough to show that the terms in G_l dominates the (mainly non-linear) terms that are in F_l but not G_l – that $|G_l| \geq |F_l - G_l|$. That part of the proof is computer-assisted – the proofs of the estimates are in this paper but computing the exact values and checking that the inequalities holds is done by a computer program using the CAPD package for rigorous interval arithmetics.

1.1 The program

This program is written in C++ and can be downloaded from <http://www.im.uj.edu.pl/MikolajZalewski/dl/delay-sin.tgz>. The rounding-mode changing code required by the interval arithmetic is system-dependent and has been checked to work on PCs (both 32-bit and 64-bit) on both Windows (compiled with cygwin) and Linux (compiled with gcc). It should also work on SPARC and Mac OS X although that has not been tested. Using other CPUs or compilers might require modifications to the rounding code.

2 Fourier coefficients

We will use the following notation: if $\{x_n\}_{n=-\infty}^{\infty}$ and $\{y_n\}_{n=-\infty}^{\infty}$ are sequences with complex values then by $x * y$ we will denote the convolution of the sequences:

$$x * y := \left\{ \sum_{k=-\infty}^{\infty} x_k y_{n-k} \right\}_{n=-\infty}^{\infty}$$

It is well known that if the sum converges the operation of convolution is associative. We will also use the notation $(\dots)_n$ for the n -th coefficient of the sequence in brackets – e.g. $(x * y)_n = \sum_{k=-\infty}^{\infty} x_k y_{n-k}$. Also:

$$x^{*k} := \underbrace{x * \dots * x}_{k \text{ times}} \quad (\text{for } k \geq 1)$$

$$(x^{*0})_n := \begin{cases} 1 & \text{if } n = 0 \\ 0 & \text{if } n \neq 0 \end{cases}$$

Of course the sum in the definition of the convolution may be not converging so we will limit our attention to a domain where the sum will be always convergent. The following lemma holds:

Lemma 1. *If x and y is such that $\exists \alpha \geq 2, \beta_1, \beta_2 > 0 : \forall n : |x_n| \leq \frac{\beta_1}{(|n|+1)^\alpha}, |y_n| \leq \frac{\beta_2}{(|n|+1)^\alpha}$, then $(x * y)_n$ is convergent for each n and $|(x * y)_n| \leq C \frac{\beta_1 \beta_2}{(|n|+1)^\alpha}$ where $C = \frac{2(2\alpha+1)}{\alpha-1}$*

Proof: Let's assume $n \geq 0$ and let's estimate $|(x * y)_n|$:

$$\left| \sum_{k=-\infty}^{\infty} x_k y_{n-k} \right| \leq \left| \sum_{k=0}^{\infty} x_{-k} y_{n+k} \right| + \left| \sum_{k=1}^{n-1} x_k y_{n-k} \right| + \left| \sum_{k=0}^{\infty} x_{n+k} y_{-k} \right|$$

$$\begin{aligned} \left| \sum_{k=1}^{n-1} x_k y_{n-k} \right| &\leq \beta_1 \beta_2 \sum_{k=1}^{n-1} \frac{1}{(k+1)^\alpha} \frac{1}{(n-k+1)^\alpha} \\ &\leq 2\beta_1 \beta_2 \sum_{1 \leq k \leq \frac{n}{2}} \frac{1}{(k+1)^\alpha} \frac{1}{(n-k+1)^\alpha} \\ &\leq 2\beta_1 \beta_2 \sum_{1 \leq k \leq \frac{n}{2}} \frac{1}{(k+1)^\alpha} \frac{1}{((n/2)+1)^\alpha} \\ &\leq 2 \frac{\beta_1 \beta_2 2^\alpha}{(n+2)^\alpha} \sum_{1 \leq k \leq \frac{n}{2}} \frac{1}{(k+1)^\alpha} \\ &\leq 2 \frac{\beta_1 \beta_2 2^\alpha}{(n+2)^\alpha} \sum_{k \geq 1} \frac{1}{(k+1)^\alpha} \end{aligned}$$

$$\begin{aligned} \left| \sum_{k=0}^{\infty} x_{-k} y_{n+k} \right| &\leq \beta_1 \beta_2 \sum_{k=0}^{\infty} \frac{1}{(k+1)^\alpha} \frac{1}{(n+k)^{\alpha+1}} \\ &\leq \frac{\beta_1 \beta_2}{(n+1)^\alpha} \sum_{k \geq 0} \frac{1}{(k+1)^\alpha} \end{aligned}$$

Analogically:

$$\left| \sum_{k=0}^{\infty} x_{n+k} y_{-k} \right| \leq \frac{\beta_1 \beta_2}{(n+1)^\alpha} \sum_{k \geq 0} \frac{1}{(k+1)^\alpha}$$

The sums $\sum_{k=k_0}^{\infty} \frac{1}{k^\alpha+1}$ can be estimated by integrals:

$$\begin{aligned} \sum_{k=0}^{\infty} \frac{1}{(k+1)^\alpha} &\leq 1 + \int_0^{\infty} \frac{1}{(x+1)^\alpha} dx = \frac{\alpha}{\alpha-1} \\ \sum_{k=1}^{\infty} \frac{1}{(k+1)^\alpha} &\leq \frac{1}{2^\alpha} + \int_1^{\infty} \frac{1}{(x+1)^\alpha} dx = \frac{\alpha+1}{2^\alpha(\alpha-1)} \end{aligned}$$

From that we obtain:

$$\left| \sum_{k=-\infty}^{\infty} x_k y_{n-k} \right| \leq \frac{\beta_1 \beta_2}{(n+1)^\alpha} \left[2^{\alpha+1} \frac{\alpha+1}{2^\alpha(\alpha-1)} + 2 \frac{\alpha}{\alpha-1} \right] \leq \frac{\beta_1 \beta_2}{(n+1)^\alpha} \frac{2(2\alpha+1)}{\alpha-1}$$

The result for $n < 0$ can be obtained by analogous estimations or by taking a sequences $\tilde{x}, \tilde{y}$: $\tilde{x}_n := x_{-n}$, $\tilde{y}_n := y_{-n}$ and applying for them the result for $n > 0$. □

With one exception we will use this lemma for $\alpha = 2$. For $\alpha = 2$ we have $C = 10$.

Observation 1. *If x is such that $\exists \alpha \geq 2, \beta > 0 : \forall n : |x_n| \leq \frac{\beta}{(|n|+1)^\alpha}$ then $|(x^{*k})_n| \leq C^{k-1} \frac{\beta^k}{(|n|+1)^\alpha}$ where $C = \frac{2(2\alpha+1)}{\alpha-1}$*

Lemma 2. *If x and y are such that $x_{-n} = \overline{x_n}$, $y_{-n} = \overline{y_n}$ then $(x * y)_{-n} = \overline{(x * y)_n}$*

Thus we have that if $x \in X_{\beta_1}$ and $y \in X_{\beta_2}$ then $x * y \in X_{C\beta_1\beta_2}$ (where X_β was defined in the introduction and C is from Lemma 1). It will be also useful to define a set of sequences from any X_β :

$$X := \bigcup_{\beta > 0} X_\beta$$

We will limit ourselves to the sequences in X . For them we have:

Lemma 3. *Let $c \in X$. Then $\sum_{n=-\infty}^{\infty} c_n e^{int}$ converges to a real-valued continuous function*

Proof: The functions $\sum_{n=-N}^N c_n e^{int}$ are continuous and real-valued as $c_n = \overline{c_{-n}} \Rightarrow c_n e^{int} + c_{-n} e^{-int} \in \mathbb{R}$. They uniformly convergent as $\sum_{n=-\infty}^{\infty} |c_n e^{int}| \leq \sum_{n=-\infty}^{\infty} \frac{\beta}{(|n|+1)^2} < \infty$. Thus the limit is also real-valued and continuous. $\square$

Let's note that $c \in X$ doesn't guarantee that the function is $\mathcal{C}^1$:

We use the convolution because of this well known fact:

Lemma 4. *If $c \in X$ are the Fourier coefficients of $x(t)$, $d \in X$ are the coefficients of $y(t)$ then the Fourier coefficients of $x(t) \cdot y(t)$ are $c * d$. As a consequence the coefficients of $x^n(t)$ are c^{*n} .*

Now we can write the equation (2) on the Fourier coefficients.

Theorem 1. *Let τ be fixed and $x(t) : \mathbb{R} \rightarrow \mathbb{R}$ be a 2π -periodic function with the Fourier coefficients $c \in X$. Then:*

(i) *for each $n \in \mathbb{Z}$ the sum on the right-hand side of the following equation converges:*

$$inc_n = -\frac{K}{\tau} \sum_{k=0}^{\infty} \frac{(-1)^k}{(2k+1)!} (c^{*(2k+1)})_n e^{-in\tau} \quad (3)$$

(ii) *$x(t)$ is a 2π -periodic solution of (2) $\Leftrightarrow c$ satisfies the equations (3).*

To prove the theorem we will need some lemmas. As it has been already mentioned that $c \in X$ doesn't imply that $x(t)$ is $\mathcal{C}^1$. However if c is the solution of equation (3) then we have the following lemma that will allow us to show that $x(t)$ is $\mathcal{C}^1$ ($x(t)$ is even $\mathcal{C}^\infty$):

Lemma 5. *If c satisfies equation (3) and $\beta > 0, \alpha \geq 2$ are such that $\forall n : |c_n| \leq \frac{\beta}{(|n|+1)^\alpha}$ then $\exists \beta' : \forall n : |c_n| \leq \frac{\beta'}{(|n|+1)^{\alpha+1}}$*

Proof: Let $C := \frac{2(2\alpha+1)}{\alpha-1}$. We have that:

$$\begin{aligned} \left| -\frac{K}{\tau} \sum_{k=0}^{\infty} \frac{(-1)^k}{(2k+1)!} (c^{*(2k+1)})_n e^{-in\tau} \right| &\leq \frac{K}{\tau} \sum_{k=0}^{\infty} \frac{1}{(2k+1)!} | (c^{*(2k+1)})_n | \\ &\leq \frac{K}{\tau} \sum_{k=0}^{\infty} \frac{1}{(2k+1)!} C^{2k} \frac{\beta^{2k+1}}{(|n|+1)^\alpha} \\ &= \frac{K}{\tau} \frac{1}{C(|n|+1)^\alpha} \sinh(C\beta) \end{aligned} \quad (4)$$

We also have $|inc_n| = |n| |c_n|$. The two sides of the equation must be equal hence we obtain:

$$\frac{K}{\tau} \frac{1}{C(|n|+1)^\alpha} \sinh(C\beta) \geq |n| |c_n|$$

Thus for $n \neq 0$ the $\beta' = \frac{2K}{C\tau} \sinh(C\beta)$ satisfies the assertion. If it is not satisfied for $n = 0$ we can increase the β' . $\square$

Increasing the α is important as we have:

Lemma 6. *Let c be a sequence of complex values satisfying $|c_n| \leq \frac{\beta}{(|n|+1)^3}$ for some β and $c_n = \overline{c_{-n}}$. Then the sequence c is a sequence of Fourier coefficients of a real-valued $\mathcal{C}^1$ function $x(t)$.*

Proof: The sequences $\sum_{k=-n}^n c_k e^{ikt}$ and $\sum_{k=-n}^n ikc_k e^{ikt}$ are real-valued, the second is the derivative of the first one and are uniformly convergent as $n \rightarrow \infty$. Hence both converge to continuous function and the second is the derivative of the first one. Thus $x(t) = \sum_{k=-\infty}^{\infty} c_k e^{ikt}$ is $\mathcal{C}^1$. $\square$

Proof of the Theorem 1:

Ad (i): The convolutions are convergent because $x \in X$. From the equation (4) from the proof of Lemma 5 we have that if $x \in X_\beta$ then $\sum_{k=0}^{\infty} \left| \frac{(-1)^k}{(2k+1)!} (c^{*(2k+1)})_n e^{-in\tau} \right| \leq \frac{1}{C(|n|+1)^2} \sinh(C\beta) < \infty$

Ad (ii): Implication $\Rightarrow$: It's enough to show that $\{inc_n\}_{n=-\infty}^{\infty}$ are the Fourier coefficients of $x'(t)$ while $-\frac{K}{\tau} \sum_{k=0}^{\infty} \frac{(-1)^k}{(2k+1)!} (c^{*(2k+1)})_n e^{-in\tau}$ are the Fourier coefficients of $-\frac{K}{\tau} \sin x(t - \tau)$.

The first part can be obtained by integrating $\int_0^{2\pi} x'(t) e^{ikt} dt$ by parts.

As for the second we have that $-\frac{K}{\tau} \sum_{k=0}^N \frac{(-1)^k}{(2k+1)!} (x(t - \tau))^{2k+1} \rightarrow -\frac{K}{\tau} \sin x(t - \tau)$ as $N \rightarrow \infty$. The Fourier coefficients of $x(t - \tau)$ are equal to $d := \{c_n e^{-in\tau}\}_{n=-\infty}^{\infty}$. Hence the coefficients of $x(t - \tau)^{2k+1}$ are $d^{*(2k+1)}$ which is equal to $c^{*(2k+1)} e^{-in\tau}$.

Hence we have that the Fourier coefficients of $-\frac{K}{\tau} \sum_{k=0}^N \frac{(-1)^k}{(2k+1)!} (x(t - \tau))^{2k+1}$ are equal to $\left\{ -\frac{K}{\tau} \sum_{k=0}^N \frac{(-1)^k}{(2k+1)!} (c^{*(2k+1)})_n e^{-in\tau} \right\}_{n=-\infty}^{\infty}$. It has been shown that this sequence is convergent as $N \rightarrow \infty$ thus the n -th coefficient of $-\frac{K}{\tau} \sin x(t - \tau)$ is $-\frac{K}{\tau} \sum_{k=0}^{\infty} \frac{(-1)^k}{(2k+1)!} (c^{*(2k+1)})_n e^{-in\tau}$. This ends the proof of this case.

Implication $\Leftarrow$: from Lemmas 5 and 6 we obtain that $x(t)$ is a $\mathcal{C}^1$ function. We know that $-\frac{K}{\tau} \sum_{k=0}^{\infty} \frac{(-1)^k}{(2k+1)!} (c^{*(2k+1)})_n e^{-in\tau}$ are the Fourier coefficients of $-\frac{K}{\tau} \sin x(t - \tau)$. They are equal to inc_n - the Fourier coefficients of $x'(t)$. So both $-\frac{K}{\tau} \sin x(t - \tau)$ and $x'(t)$ are continuous and 2π -periodic functions with equal Fourier coefficients. Hence the functions themselves are equal and equation (2) is satisfied. $\square$

Let's define a function $F : \mathbb{R} \times X \rightarrow \mathbb{C}^{\mathbb{Z}}$ (with the product topology on $\mathbb{C}^{\mathbb{Z}}$) corresponding to the equation (3):

$$F(\tau, c) := \left\{ in\tau e^{in\tau} c_n + K \sum_{k=0}^{\infty} \frac{(-1)^k}{(2k+1)!} (c^{*(2k+1)}) \right\}_{n=-\infty}^{\infty}$$

Of course having $F(\tau, c) = 0$ is equivalent to the fact that τ, c satisfies equation (3). In the rest of the paper we will show that F has a nontrivial zero. Let's note that $F(\tau, c)_n = F(\tau, c)_{-n}$.

As we will use topological tools we will want F to be continuous. First let's note that the convolution is not continuous on the whole X (as written above we use the product topology on X) but we have:

Lemma 7. *The operation of convolution is continuous on each X_β*

Proof: Let's fix some $x^0, y^0 \in X_\beta$. Let's fix some δ and take some $x, y \in X_\beta$ from a neighborhood of (x^0, y^0) : x, y are such that $\forall |n| \leq N_\delta : |x_n - (x^0)_n|, |y_n - (y^0)_n| < \delta$ where N_δ is large enough that for $|n| > N_\delta : x, y, x^0, y^0 \in X_\beta \Rightarrow |x_n - (x^0)_n|, |y_n - (y^0)_n| < \delta$. We have that:

$$|(x^0 * y^0 - x * y)_n| \leq |(x^0 * (y^0 - y))_n| + |((x^0 - x) * y)_n|$$

$$|(x^0 * (y^0 - y))_n| \leq \sum_{j=-\infty}^{\infty} |(x^0)_j| \cdot |(y^0)_{n-j} - y_{n-j}| \leq \beta \delta \sum_{j=-\infty}^{\infty} \frac{1}{(|j| + 1)^2}$$

This tends to zero as $\delta \rightarrow 0$. After an analogous estimation for $|((x^0 - x) * y)_n|$ we have that the convolution is continuous. $\square$

Lemma 8. *The function F is continuous on each X_β .*

Proof: The convolutions are continuous on each X_β and the result of a convolution lays in a $X_{\beta'}$ for some β' so $K \sum_{k=0}^N \frac{(-1)^k}{(2k+1)!} (c^{*(2k+1)})_n$ is continuous for each N . From estimations as in equation (4) we have that for each coefficient this series converges uniformly as $N \rightarrow \infty$ hence we obtain that the limit $K \sum_{k=0}^{\infty} \frac{(-1)^k}{(2k+1)!} (c^{*(2k+1)})_n$ is continuous.

The term $in\tau e^{in\tau} c_n$ is also continuous so F is continuous. $\square$

3 Some estimates

As mentioned in the introduction we will need some estimates to show that the inequality holds in the neighborhood of $\hat{c}$. We will use two kinds of sets:

$\hat{c} + X_{\beta_2}$ that will be used to obtain finer estimates and X_{β_1} (where β_1 will be large enough to contain the whole set $\hat{c} + X_{\beta_2}$) for some more rough but simpler ones.

In this section we will only assume of $\hat{c}$ that almost all coefficients are equal to zero. By Y_l we will denote the space of the possible values of $\hat{c}$ – the set of sequences such that at most the elements $-l, \dots, l$ are non-zero:

$$Y_l := \{c \in X : \forall n : |n| > l \Rightarrow c_n = 0\}$$

First let's note two simple properties:

Lemma 9. *If $c \in Y_l$ then $c^{*k} \in Y_{kl}$*

Lemma 10. *If $x, y \in X$ then*

$$(x + y)^{*n} = \sum_{k=0}^{k=n} \binom{n}{k} x^{*k} * y^{*(n-k)}$$

To estimate $\frac{K}{p!} (x^{*p})_n$ – an element of the sum in the equation (3) – for $x \in X_\beta$ it's enough to use Lemma 1. However for the sets of the form $c + X_\beta$ we will use a more sophisticated estimation:

Lemma 11. *Let $c \in Y_l$, $p > 1$, $\beta > 0$. Then for any $x \in X_\beta$:*

$$\left| \frac{K}{p!} (c + x)_n^{*p} \right| \leq \frac{K}{p!} \left(|c_n^{*p}| + \sum_{k=0}^{p-1} \binom{p}{k} \sum_{j=-lk}^{lk} |c_j^{*k}| C^{p-k-1} \frac{\beta^{p-k}}{(|n-j|+1)^2} \right)$$

Where $C = 10$

Proof: From Lemmas 1 and 10 we have:

$$\begin{aligned} \left| \frac{K}{p!} (x + c)_n^{*p} \right| &\leq \frac{K}{p!} \sum_{k=0}^p \binom{p}{k} |(c^{*k} * x^{*(p-k)})_n| \\ &\leq \frac{K}{p!} \sum_{k=0}^p \binom{p}{k} \sum_{j=-\infty}^{\infty} |c_j^{*k}| |x_{n-j}^{*(p-k)}| \\ &\leq \frac{K}{p!} \left(|c_n^{*p}| + \sum_{k=0}^{p-1} \binom{p}{k} \sum_{j=-\infty}^{\infty} |c_j^{*k}| C^{p-k-1} \frac{\beta^{p-k}}{(|n-j|+1)^2} \right) \end{aligned}$$

We have $c \in Y_l$ so $c_j^{*k} = 0$ for $j > kl$ what ends the proof. $\square$

Of course we can use the previous lemma only for a finite number of terms. To estimate the tail we will use a weaker estimation by using a neighborhood of the second type, applying Lemma 1 and summing the geometric sequence:

Lemma 12. Let N be odd, $\beta \in [0; \frac{N}{10}]$. If $x \in X_\beta$ then

$$\left| \left(\sum_{k=\frac{N-1}{2}}^{\infty} \frac{K}{(2k+1)!} x^{*(2k+1)} \right)_n \right| \leq \frac{K \cdot (C\beta)^{N-1}}{N!(1 - (\frac{C\beta}{N})^2)} \cdot \frac{\beta}{(|n|+1)^2}$$

Where $C = 10$.

Proof: Using Lemma 1:

$$\begin{aligned} \left| \left(\sum_{k=\frac{N-1}{2}}^{\infty} \frac{K}{(2k+1)!} x^{*(2k+1)} \right)_n \right| &\leq \sum_{k=(N-1)/2}^{\infty} \frac{K}{N!N^{2k-(N-1)}} \left| (x^{*(2k+1)})_n \right| \\ &\leq \frac{K}{N!} \sum_{k=(N-1)/2}^{\infty} \frac{(C\beta)^{2k}}{N^{2k-(N-1)}} \cdot \frac{\beta}{(|n|+1)^2} \\ &= \frac{K(C\beta)^{N-1}}{N!} \left(\sum_{k=0}^{\infty} \left(\frac{C^2\beta^2}{N^2} \right)^k \right) \cdot \frac{\beta}{(|n|+1)^2} \\ &= \frac{K(C\beta)^{N-1}}{N!(1 - (\frac{C\beta}{N})^2)} \cdot \frac{\beta}{(|n|+1)^2} \end{aligned}$$

The geometric sequence is convergent because $\beta < \frac{N}{10} \Rightarrow (\frac{C\beta}{N})^2 < 1$. $\square$

From Lemma 11 we can obtain an estimation that after multiplication by $(|n|+1)^2$ is independent of n . That will be used to estimate the term in all inequalities for high n -s with one formula:

Lemma 13. Let $\beta > 0$, $c \in Y_l$, $p > 1$, $N > pl$. Then for each $x \in X_\beta$, $n > N$ we have:

$$\left| \frac{K}{p!} (x+c)_n^{*(p)} \right| \leq \frac{K}{p!} \left(\sum_{k=0}^{p-1} \binom{p}{k} \sum_{j=-lk}^{lk} |c_j^{*k}| (C\beta)^{p-k-1} \frac{(N+1)^2}{(N-j+1)^2} \right) \frac{\beta}{(|n|+1)^2}$$

Proof: Note that $n > pl \Rightarrow c_n^{*p} = 0$ and $\frac{\beta}{(|n-j|+1)^2} = \frac{(n+1)^2}{(|n-j|+1)^2} \cdot \frac{\beta}{(n+1)^2} \leq \frac{(N+1)^2}{(N-j+1)^2} \cdot \frac{\beta}{(n+1)^2}$ as $N > j$ and apply Lemma 11 $\square$

For the first terms we will need to have a better estimation than in Lemma 11 so we will regroup the terms (to understand why this regrouping helps let's compare two estimates: $|1.1 \cdot x - 1 \cdot x| \leq 1.1|x| + 1|x| = 2.1|x|$ and $|1.1 \cdot x - 1 \cdot x| = |(1.1 - 1)x| = 0.1|x|$).

Lemma 14. Let $x \in X_\beta$, $c \in Y_l$.

$$\begin{aligned} in\tau e^{in\tau} (c_n + x_n) + K \sum_{k=0}^3 \frac{(-1)^k}{(2k+1)!} (c+x)_n^{*(2k+1)} = \\ (in\tau e^{in\tau} + K) c_n - \frac{K}{3!} (c^{*3})_n + \frac{K}{5!} (c^{*5})_n - \frac{K}{7!} (c^{*7})_n + \\ + (in\tau e^{in\tau} + K) x_n + \sum_{p=1}^7 \sum_{j=-6l}^{6l} \gamma_{p,j} x_{n-j}^{*p} \end{aligned}$$

Where $\gamma_{p,j} = -\frac{K}{3!} \binom{3}{p} (c^{*(3-p)})_j + \frac{K}{5!} \binom{5}{p} (c^{*(5-p)})_j - \frac{K}{7!} \binom{7}{p} (c^{*(7-p)})_j$
and we assume $\binom{n}{k} = 0$ for $k > n$.

4 Proving the existence of the periodic solution

We will search for the orbit in the neighborhood of $\hat{c}$ and $\hat{\tau}$ defined in the introduction. Let's define the sets and boundaries on which we will operate:

$$X_1 := X_{\beta_1}$$

$$X_2 := \hat{c} + X_{\beta_2}$$

$$X_3 := \{y \in X_2 : y_1 - \hat{c}_1 \in \mathbb{R}\}$$

$$\underline{\tau} := \hat{\tau} - \Delta\tau$$

$$\bar{\tau} := \hat{\tau} + \Delta\tau$$

where $\Delta\tau = 0.000001$, $\beta_2 = 0.0000002438$, $\beta_1 = 0.766763$.

We will prove that a solution exists in the set $[\underline{\tau}; \bar{\tau}] \times X_3$. The β_1, β_2 are such that $X_3 \subset X_2 \subset X_1$.

As mentioned in the introduction we will use the Galerkin projections of F with the condition $y_1 - \hat{c}_1 \in \mathbb{R}$ added to make the zero isolated. The P_l and Q_l were defined in the introduction as the projection and immersion of the finite-dimensional space. Let's first prove the lemma stated in the introduction.

Lemma 15. *Let $\beta > 0$, $l_0 > 0$, $\underline{\tau}, \bar{\tau} \in \mathbb{R}$ be fixed. If for each $l > l_0$ there is a $c^l \in X_\beta$ and $\tau_l \in [\underline{\tau}, \bar{\tau}]$ such that $P_l F(\tau_l, c^l) = 0$ then there exist $(c^0, \tau_0) \in X_\beta \times [\underline{\tau}, \bar{\tau}]$ such that $F(\tau_0, c^0) = 0$.*

Moreover, if all the c^l are in a closed set D then $c^0 \in D$.

Proof: From the compactness of $[\underline{\tau}, \bar{\tau}] \times X_\beta$ there exists a subsequence l_k such that c^{l_k} converges to a limit c^0 and τ_{l_k} converges to τ_0 . Let's fix $n \geq 0$ and note that for $l > n$ if $P_l(F(\tau, c)) = 0$ then $F(\tau, c)_n = 0$, so from the continuity of F we have $F(\tau, c^0)_n = \lim_{k \rightarrow \infty} F(\tau_{l_k}, c^{l_k})_n = \lim_{k \rightarrow \infty} \overline{P_{l_k} F(\tau_{l_k}, c^{l_k})_n} = 0$. For $n < 0$ we have $F(\tau_0, c^0)_n = \lim_{k \rightarrow \infty} F(\tau_{l_k}, c^{l_k})_n = \lim_{k \rightarrow \infty} F(\tau_{l_k}, c^{l_k})_{-n} = 0$. Thus $F(\tau_0, c^0) = 0$.

The last assertion is obvious. $\square$

Before defining the homotopy let's introduce some auxiliary notations. Let's denote: $f_n(\tau) := in\tau e^{in\tau}$. By L_n we will denote the linear part of f_n :

$L_n(\tau) := (ine^{in\tau_0} - n^2e^{in\tau_0})(\tau - \tau_0)$. By r_n we will denote the non-linear part:
 $r_n(\tau) := f_n(\tau) - f_n(\tau_0) - L_n(\tau)$.

We will define the homotopy on the whole infinite-dimensional space:
 $H : [0; 1] \times [\underline{\tau}; \overline{\tau}] \times X_3 \rightarrow \mathbb{C}^{\mathbb{Z}}$. We will use only the Galerkin projections of this homotopy to deform F_l (that is the Galerkin projection of F – see introduction) but to prove that there are no zeros on the boundaries it's a bit more convenient to define the homotopy on the whole space. As written in the introduction the homotopy is a linear deformation of F to a nearly linear function G :

$$H(h, \tau, x) := hF(\tau, \hat{c} + x) + (1 - h)G(\tau, x)$$

where G on the n -th coefficient is equal to:

$$G(\tau, x)_n := \begin{cases} (f_n(\tau) + K) x_n & \text{for } n \neq \pm 1 \\ (f_n(\tau_0) + K + \gamma_{1,0} + \gamma_{1,2n}) x_n + L(\tau) \hat{c}_n & \text{for } n = \pm 1 \end{cases}$$

Where $\gamma_{p,j}$ is from Lemma 14.

The G is not strictly a linearization of F as it doesn't contain all the linear terms and is not linear with respect to τ . However it contains the most important terms – the rest will be shown to be small compared to them. Let's note that for $n = \pm 1$ we have $x_{n-2n} = \overline{x_n} = x_n$ (because $x_{\pm 1} \in \mathbb{R}$) so the term with $\gamma_{1,2n}$ in F will be a linear term with respect to x_n .

One can write explicit formulas for H . For $n \neq \pm 1$ we have:

$$H(h, \tau, x)_n := (f_n(\tau) + K) x_n + \\ + h \left((f_n(\tau) + K) \hat{c}_n + \sum_{k=1}^{\infty} \frac{K}{(2k+1)!} ((x + \hat{c})^{*(2k+1)})_n \right)$$

And for $n = \pm 1$:

$$H(t, \tau, x)_n := (f_n(\tau_0) + K + \gamma_{1,0} + \gamma_{1,2n}) x_n + L_n(\tau) \hat{c}_n + \\ + h \left((f_n(\tau_0) + K) \hat{c}_n - \frac{K}{3!} (\hat{c}^{*3})_n + \frac{K}{5!} (\hat{c}^{*5})_n - \frac{K}{7!} (\hat{c}^{*7})_n \right) + \\ + h \sum_{-6.5 \leq j \leq 6.5, j \neq 0, 2n} \gamma_{1,j} x_{n-j} + h \sum_{p=2}^7 \sum_{j=-6.5}^{6.5} \gamma_{p,j} x_{n-j}^{*p} + \\ + h r_n(\tau) \hat{c}_n + h(f_n(\tau) - f_n(\tau_0)) x_n + h R(\hat{c} + x)$$

Where R is a short notation for:

$$R(y) := \sum_{k=4}^{\infty} \frac{K}{(2k+1)!} y^{*(2k+1)}$$

To show that any Galerkin projection of H doesn't have a zero on the boundary of $[\underline{\tau}; \overline{\tau}] \times P_l(X_3)$ it's enough to show that H doesn't have a zero on the boundary. More exactly:

Theorem 2. Let $x \in X_3$, $\tau \in [\underline{\tau}, \bar{\tau}]$. Let x be such that $\exists n : |x_n| = \frac{\beta_2}{(|n|+1)^2}$ or let $\tau \in \{\underline{\tau}, \bar{\tau}\}$. Then $\forall h \in [0; 1] : H(h, \tau, x) \neq 0$.

Proof: The proof is computer assisted – the calculations are done by the program. The computation is as follow.

First let's note that if for $n < 0$ we have $|x_n| = \frac{\beta_2}{(|n|+1)^2}$ then this is also true for $-n$. Thus it's enough to consider $n \geq 0$ (and τ). We do it in two steps:

1. Let's assume that $|x_n| = \frac{\beta_2}{(|n|+1)^2}$ for some $n \geq 0$, $n \neq 1$. We will prove that:

$$|(f_n(\tau) + K) x_n| > \left| (f_n(\tau) + K) \hat{c}_n + \sum_{k=1}^{\infty} \frac{K}{(2k+1)!} ((x + \hat{c})^{*(2k+1)})_n \right|$$

As the value of $H(h, \tau, x)_n$ is the difference of these two terms (with the second one multiplied by a $h < 1$), it will follow that $H(h, \tau, x)_n \neq 0$ thus $H(h, \tau, x) \neq 0$. The LHS will be estimated by:

$$|(f_n(\tau) + K) x_n| \geq ||f_n(\tau)| - K| \frac{\beta_2}{(|n| + 1)^2} \quad (5)$$

and $|f_n(\tau)| = n|\tau|$ is an interval in the interval arithmetic.

The terms of RHS are estimated depending on n . For $n < 225$ we can estimate each n separately. For $n \geq 225$ we will want to check all the inequalities in some finite computations.

Let's note that for $n \geq 255$ we have $(f_n(\tau) + K) \hat{c}_n = 0$ as $\hat{c}_n = 0$. In the sum $\sum_{k=1}^{\infty} \frac{K}{(2k+1)!} ((x + \hat{c})^{*(2k+1)})_n$ we will estimate the terms $k = 1, \dots, 22$ from Lemma 13 while the term $k > 22$ from Lemma 12. Now if we multiply the estimates for both LHS and RHS by $(|n| + 1)^2$ they will be independent of n . Hence we will check all the $n \geq 225$ by checking only one inequality.

For $n < 225$ we have to use some more sophisticated estimations. Let's use Lemma 14 to group the terms:

$$\begin{aligned} & \left| (f_n(\tau) + K) \hat{c}_n + \sum_{k=1}^{\infty} \frac{K}{(2k+1)!} ((x + \hat{c})^{*(2k+1)})_n \right| \\ = & \left| ((f_n(\tau) + K) \hat{c}_n - \frac{K}{3!} (\hat{c}^{*3})_n + \frac{K}{5!} (\hat{c}^{*5})_n - \frac{K}{7!} (\hat{c}^{*7})_n) + \right. \\ & \left. \sum_{p=1}^7 \sum_{j=-6l}^{6l} \gamma_{p,j} x_{n-j}^{*k} + R(\hat{c} + x) \right| \\ \leq & \left| (f_n(\tau) + K) \hat{c}_n - \frac{K}{3!} (\hat{c}^{*3})_n + \frac{K}{5!} (\hat{c}^{*5})_n - \frac{K}{7!} (\hat{c}^{*7})_n \right| + \\ & \left| \sum_{p=1}^7 \sum_{j=-6l}^{6l} \gamma_{p,j} x_{n-j}^{*k} \right| + |R(\hat{c} + x)| \\ \leq & |(f_n(\tau) - f_n(\tau_0)) \hat{c}_n| + \\ & + \left| (f_n(\tau_0) + K) \hat{c}_n - \frac{K}{3!} (\hat{c}^{*3})_n + \frac{K}{5!} (\hat{c}^{*5})_n - \frac{K}{7!} (\hat{c}^{*7})_n \right| + \\ & + \sum_{p=1}^7 \sum_{j=-6l}^{6l} |\gamma_{p,j}| \frac{\beta_2 (10\beta_2)^{p-1}}{(|n-j|+1)^2} \\ & + \sum_{k=4}^{22} \left| ((x + \hat{c})^{*(2k+1)})_n \right| + \left| \sum_{k=23}^{\infty} ((x + \hat{c})^{*(2k+1)})_n \right| \end{aligned}$$

Table 1: Estimates of LHS and RHS for small n

n	LHS	RHS
0	$4.1792 \cdot 10^{-7}$	$0.1687 \cdot 10^{-7}$
2	$0.4474 \cdot 10^{-7}$	$0.0958 \cdot 10^{-7}$
3	$0.5080 \cdot 10^{-7}$	$0.0827 \cdot 10^{-7}$
4	$0.4893 \cdot 10^{-7}$	$0.0160 \cdot 10^{-7}$
5	$0.4537 \cdot 10^{-7}$	$0.0118 \cdot 10^{-7}$
6	$0.4171 \cdot 10^{-7}$	$0.0070 \cdot 10^{-7}$
7	$0.3834 \cdot 10^{-7}$	$0.3465 \cdot 10^{-7}$
8	$0.3536 \cdot 10^{-7}$	$0.0040 \cdot 10^{-7}$
9	$0.3274 \cdot 10^{-7}$	$0.0036 \cdot 10^{-7}$

The estimates obtained by the program. The LHS is closest to the RHS for $n = 7$ because we have artificially chosen $\hat{c}_7 = 0$. That makes the term $|(f_n(\tau) + K)\hat{c}_n - \frac{K}{3!}(\hat{c}^{*3})_n + \frac{K}{5!}(\hat{c}^{*5})_n - \frac{K}{7!}(\hat{c}^{*7})_n|$ for $n = 7$ equal to approximately $0.3411 \cdot 10^{-7}$.

in the first term of RHS we will use the estimation:

$$\begin{aligned}
|(f_n(\tau) - f_n(\tau_0))| &\leq \max_{\tau \in [\underline{\tau}, \bar{\tau}]} |f'(\tau)(\tau - \tau_0)| \leq \\
&\leq \max_{\tau \in [\underline{\tau}, \bar{\tau}]} |ine^{in\tau} - n^2\tau e^{in\tau}| \cdot |\Delta\tau| \leq \\
&\leq \max_{\tau \in [\underline{\tau}, \bar{\tau}]} (n + n^2\tau) |\Delta\tau|
\end{aligned} \tag{6}$$

The fact that it grows quickly with n is irrelevant as for $n > 5$ we have $\hat{c}_n = 0$.

The $|(f_n(\tau_0) + K)\hat{c}_n - \frac{K}{3!}(\hat{c}^{*3})_n + \frac{K}{5!}(\hat{c}^{*5})_n - \frac{K}{7!}(\hat{c}^{*7})_n|$ can be directly computed and is small as it is the numerical solution that is close to zero.

The $\sum_{p=1}^7 \sum_{j=-6l}^{6l} |\gamma_{p,j}| \frac{\beta_2(10\beta_2)^{p-1}}{(n-j+1)^2}$ can be directly computed and is small for $p > 1$ as $10\beta_2$ is small. If the terms for $p = 1$ wouldn't be small we could try to diagonalize linear part. But there is no such need – we check that the inequality holds. We estimate $|((x + \hat{c})^{*(2k+1)})_n|$ for $k \in \{4, \dots, 22\}$ from Lemma 11. To estimate $|\sum_{k=23}^{\infty} ((x + \hat{c})^{*(2k+1)})_n|$ we use Lemma 12. In both estimates we have $\frac{1}{(2k+1)!}$ with k large enough to make the result small.

Thus we see that unless the $\gamma_{1,\cdot}$ are big, the estimation of RHS should be small. The program computes the estimation for LHS and RHS, compares them and for each n in $0, 2, 3, 4, \dots, 224$ the inequality to hold. This ends the case for $n \neq 1$.

2. We have two cases left: $\tau \in \{\underline{\tau}, \bar{\tau}\}$ and $|x_1| = \frac{\beta_2}{4}$. We have two variables left but only one equation – the equation for $n = 1$. However this

is a complex equation and the variables are real so we will be able to show that in both cases $H(\tau, x) \neq 0$.

Let's denote:

$$\begin{aligned} L_\tau &:= L_1(\tau)\hat{c}_1 \\ L_x &:= (f_n(\tau_0) + K + \gamma_{1,0} + \gamma_{1,2})x_1 \\ N &:= (f_n(\tau_0) + K)\hat{c}_n - \frac{K}{3!}(\hat{c}^{*3})_n + \frac{K}{5!}(\hat{c}^{*5})_n - \frac{K}{7!}(\hat{c}^{*7})_n + \\ &\quad + \sum_{-6l \leq j \leq 6l, j \neq 0, 2} \gamma_{p,j} x_{n-j}^{*k} + \sum_{p=2}^7 \sum_{j=-6l}^{6l} \gamma_{p,j} x_{n-j}^{*k} + \\ &\quad + R(c) + r_n(\tau)\hat{c}_n + (f(\tau) - f(\tau_0))x_n \end{aligned}$$

The L_τ, L_x, N depend on τ, x but to make the notation short we won't write it. They are chosen such that $G(\tau, x)_1 = L_\tau + L_x$ and $F(\tau, x)_1 = L_\tau + L_x + N$ and $H(h, \tau, x)_1 = L_\tau + L_x + hN$. If $\tau \in [\underline{\tau}, \bar{\tau}]$ then we show that:

$$|L_\tau| > |L_x| + |N|$$

The $|L_\tau|$ is estimated as:

$$|L_1(\tau)\hat{c}_1| = |in - n^2| \cdot \Delta\tau \cdot |\hat{c}_n| \Bigg|_{n=1} = \sqrt{2}\Delta\tau \cdot |\hat{c}_1|$$

We estimate the terms in $|N|$ as for $n \neq 1$ with the exception for the $|(f(\tau) - f(\tau_0))x_n|$ and $|r_n(\tau)\hat{c}_n|$ that were not present in the previous case. The the first one we use the inequality (6). For the second we use the estimation:

$$\begin{aligned} |r_n(\tau)\hat{c}_n| &\leq \sup_{\tau \in [\underline{\tau}, \bar{\tau}]} \left| \frac{1}{2} f_1''(\tau)(\tau - \tau_0)^2 \right| \cdot |\hat{c}_n| \\ &= \frac{|\hat{c}_n|}{2} \sup_{\tau \in [\underline{\tau}, \bar{\tau}]} |(2n^2 + in^3\tau)e^{in\tau}| \\ &= \frac{|\hat{c}_n|}{2} \sup_{\tau \in [\underline{\tau}, \bar{\tau}]} |2 + i\tau| \end{aligned}$$

Having all these estimates our program checks that the inequality is satisfied.

In the case $|x_1| = \frac{\beta_2}{4}$ the inequality $|L_x| > |L_\tau| + |N|$ is of course false. To prove that there is no zero will need to use the fact that $\tau \in \mathbb{R}$ and $x_1 \in \mathbb{R}$. On the figure 1 we have a sketch how the sets L_τ and $L_x + N$ looks like – we see that they shouldn't intersect.

Formally we will show that $\{\tan \arg(L_\tau) : (\tau, x) \in [\underline{\tau}, \bar{\tau}] \times X_3, x_1 = \pm \frac{\beta_2}{4}\} \cap \{\tan \arg(L_x + N) : (\tau, x) \in [\underline{\tau}, \bar{\tau}] \times X_3, x_1 = \pm \frac{\beta_2}{4}\} = \emptyset$ (where $\arg$ is the complex number argument). Of course this implies that the sum of these terms is different from zero.




Figure 1: The possible values of L_τ and $L_x + N$

The $\arg(L_\tau)$ is easy to compute as this is $\tan \arg((ie^{i\tau_0} - e^{i\tau_0})c_1)$ and $\tan \arg z$ for a complex number z can be computed as $\frac{\Im z}{\Re z}$. To estimate the other set we will use the estimation for N from the previous point – let's assume that $|N| \leq \lambda$ for some $\lambda > 0$. We have that $L_x + N \in \pm(f_n(\tau) + K + \gamma_{1,0} + \gamma_{1,2})\frac{\beta_2}{4} + [-\lambda; \lambda] + [-\lambda; \lambda]i$ and using the interval arithmetics we can find $\tan \arg(L_x + N)$. The program checks that these two sets are disjoint and this ends the proof of the theorem. $\square$

To finish the proof of the existence of the orbit let's define a second homotopy.

$$H^L(h, \tau, x) := \begin{cases} (f_n(\tau_0 + h(\tau - \tau_0)) + K) x_n & \text{for } n \neq \pm 1 \\ (f_n(\tau_0) + K + \gamma_{1,0} + \gamma_{1,2n}) x_n + L(\tau)c_1 & \text{for } n = \pm 1 \end{cases}$$

It deforms G into:

$$G^L(h, \tau, x) := \begin{cases} (f_n(\tau_0) + K) x_n & \text{for } n \neq \pm 1 \\ (f_n(\tau_0) + K + \gamma_{1,0} + \gamma_{1,2n}) x_n + L(\tau)c_1 & \text{for } n = \pm 1 \end{cases}$$

Lemma 16. *If $(\tau, x) \in [\underline{\tau}, \bar{\tau}] \times X_3$ such that $\tau \in \{\underline{\tau}, \bar{\tau}\}$ or $\exists n : |x_n| = \frac{\beta_2}{(|n|+1)^2}$ then $H^L(h, x, \tau) \neq 0$*

Proof: Let h, x, τ be such that $H^L(h, x, \tau) = 0$.

Let $n \neq \pm 1$. The $\tau_0 + h(\tau - \tau_0) \in [\underline{\tau}, \bar{\tau}]$ thus $|f_n(\tau_0 + h(\tau - \tau_0)) + K|$ can be estimated as in equation (5). For each such value we have proven that it is strictly greater than an $\text{RHS} \geq 0$. Hence $|f_n(\tau_0 + h(\tau - \tau_0)) + K| > 0$ and $H^L(h, \tau, x) = 0 \Rightarrow x_n = 0$.

But if $x_n = 0$ for each $n \neq \pm 1$ then $H^L(h, \tau, x) = G(\tau, x) = H(0, \tau, x)$ and H has no zeros on the boundary. $\square$

Observation 2. *The Galerkin projection of H : i.e. $H_l(\tau, y_0, \dots, y_l) := P_l H(\tau, Q_l(y_0, \dots, y_l))$ (for $(y_0, \dots, y_l) \in \mathbb{R} \times \mathbb{R} \times \mathbb{C}^{n-1}$) is a homotopy from $F^l(\tau, c + \cdot)$ to the projection of G . Analogically the projection of H^L is a homotopy from G to G^L . There are no zeros on the boundaries for these homotopies.*

Observation 3. *The degree $\deg(F_l, [\underline{\tau}, \bar{\tau}] \times P_l(X_3), 0)$ is well defined.*

Proof: The homotopy H_l for $h = 1$ have no zeros on the boundary and this is F_l . $\square$

Lemma 17. $\deg(F_l, [\underline{\tau}, \bar{\tau}] \times P_l(X_3), 0) \neq 0$

Proof: We know that the degree of F_l is equal to the degree of the Galerkin projection of G^L – let's denote it by G_l^L . The G_l^L is a linear function. If the determinant of the differential was zero then there would be a zero on the boundary of any neighborhood of $(\tau_0, 0)$ so the determinant is non-zero. That means that the degree is ± 1 thus it is non-zero. $\square$

Theorem 3. *The equation (2) has a periodic solution for some $\tau \in [\underline{\tau}, \bar{\tau}]$ whose Fourier coefficients are in the set X_3 .*

Proof: From the Lemma 17 we know that the local Brower degree is non-zero thus each F_l has a zero in X_3 . From Lemma 15 we have that F has a zero in X_3 . From Theorem 1 we obtain that in X_3 there is a solution of the equation. $\square$

5 Conclusions

In this paper I was able to rigorously prove the existence of a periodic orbit for $K = 1.6$. I was not able to show that the period is 4 even if the numerical simulations suggests that – in the proof we need the τ as a variable for the image and the domain to have the same dimension.

The value $K = 1.6$ has been chosen because it's easiest to find the $\hat{c}_n$ values for an attracting orbit. However using the Newton method it should

be possible to find an approximated orbit which is not attracting (from the numerical simulations it seems to happen for $K > 5.11$). For larger K the values of $\hat{c}_n$ may not decay as fast so we may need to diagonalize the first coefficients of F_l as mentioned in the proof.

Of course $\sin$ is periodic so there exist infinitely many such orbit that differs by $2k\pi$ ($k \in \mathbb{Z}$). As mentioned for $K > 5.11$ it seems that these periodic orbits stop to be attracting and orbits from numerical simulations jump $\pm 2\pi$ from one periodic orbit to another. This suggests heteroclinic connections and chaos but proving it would require some new ideas.

6 Acknowledgements

I would like to thank Piotr Zgliczyński for suggesting the topic and helpful discussions.

References

- [Cesari 64] L. Cesari, Functional analysis and Galerkin methods, *Michigan Math. J.* **11** (1964) 383-414
- [Wischert et al 94] W. Wischert, W. Wunderlin, A. Pelster, M. Olivier, and J. Grosblum *Delay-induced instabilities in nonlinear feedback systems*, Physical Review E **49** 203-219 (1994)
- [Smoller 83] J. Smoller, *Shock Waves and Reaction-Diffusion Equations* Springer, 1983.
- [Zgliczyński 01] P. Zgliczyński, K. Mischaikow *Rigorous Numerics for Partial Differential Equations: The Kuramoto-Sivashinsky Equation*, Found. Comput. Math **1** 255-288 (2001)
- [Zgliczyński 04] P. Zgliczyński *Rigorous Numerics for Dissipative Partial Differential Equations II. Periodic Orbit for the Kuramoto-Sivashinsky Equation*, Found. Comput. Math 157-195 (2004)